

Pediatric Dosing Chart

Dosage for children from your healthcare professional



DO NOT use any other dosing device.

Use products **ONLY** as directed. Always read and follow the label on all MYLICON® products. Keep out of reach of children.

Children's MYLICON® Antacid Liquid



Available in: **Bubblegum**

AGE	under 2 yrs	2 to 5 yrs	6 to 11 yrs
WEIGHT	under 24 lbs.	24 - 47 lbs.	48 - 95 lbs.
DOSE	Ask a doctor	5 mL 	10 mL 

DOSING: 5 mL - not more than 15 mL in 24 hours.
10 mL - not more than 30 mL in 24 hours.
Antacids may interact with certain drugs. Ask a doctor or pharmacist before use if your child is presently taking a prescription drug.

Infants' MYLICON® Gas Relief Drops



Available in: **Original, Dye Free**



AGE	under 2 yrs	over 2 yrs
WEIGHT	under 24 lbs.	over 24 lbs.
DOSE	0.3 mL	0.6 mL

Can be used at every feeding, up to 12 times a day.
Dosage can also be mixed with 1 oz cool water, infant formula, or other suitable liquids.

Children's MYLICON® Antacid Chewable Tablets



Available in: **Cherry**

AGE	under 2 yrs	2 to 5 yrs	6 to 11 yrs
WEIGHT	under 24 lbs.	24 - 47 lbs.	48 - 95 lbs.
DOSE	Ask a doctor	1 tablet 	2 tablets 

DOSING: 1 tablet - not more than 3 tablets in 24 hours.
2 tablets - not more than 6 tablets in 24 hours.
Antacids may interact with certain drugs. Ask a doctor or pharmacist before use if your child is presently taking a prescription drug.

Infants' MYLICON® Liquid Probiotic



Available in: **Newborns+**

AGE	Newborns
WEIGHT	—
DOSE	6 drops a day

DOSING: Do not exceed 6 drops in 24 hours.
If your child was born prematurely or has immune problems, consult your doctor prior to use.

Today's date: _____

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

© Johnson & Johnson Consumer Inc. 2024 · MYC-1134

ACTIVE INGREDIENTS: Calcium carbonate 400 mg, simethicone emulsion USP 133.3 mg/simethicone 40 mg (Children's MYLICON® Antacid Liquid,) Simethicone 20 mg (Infants' MYLICON® Gas Relief Drops,) Calcium carbonate 400 mg, simethicone 40 mg (Children's MYLICON® Antacid Chewable Tablets,) Bifidobacterium animalis subsp., lactis 1 billion CFU (Infants' MYLICON® Liquid Probiotic.)