

Pediatric Dosing Chart



Dosage for children from your healthcare professional

Use **ONLY** enclosed dosing cup specifically designed for use with this product. **DO NOT** use any other dosing device. Use products **ONLY** as directed. Always read and follow the label on all SUDAFED® products.

Children's SUDAFED® Nasal Decongestant

Available in: **Grape**





AGE	under 4 yrs	4 to 5 yrs	6 to 11 yrs
DOSE	Do not use	5 mL 	10 mL 

Active Ingredient: Pseudoephedrine HCL
Dose every 4 to 6 hours.
DO NOT give more than 4 doses in 24 hours.
Only use product as needed or as symptoms persist.

Children's SUDAFED® PE Nasal Decongestant

Available in: **Berry**






AGE	under 4 yrs	4 to 5 yrs	6 to 11 yrs
DOSE	Do not use	5 mL 	10 mL 

Active Ingredient: Phenylephrine HCL
Dose every 4 hours.
DO NOT give more than 6 doses in 24 hours.
Only use product as needed or as symptoms persist.

Children's SUDAFED® PE Cold + Cough

Available in: **Grape**



AGE	under 4 yrs	4 to 5 yrs	6 to 11 yrs
DOSE	Do not use	5 mL 	10 mL 

Active Ingredients: Phenylephrine HCL and Dextromethorphan
Dose every 4 hours.
DO NOT give more than 6 doses in 24 hours.
Only use product as needed or as symptoms persist.

Today's date: _____

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

© Johnson & Johnson Consumer Inc. 2024