ALZHEIMER'S DISEASE: TIME FOR CHANGE NOW

There is a need to **revolutionise the management of Alzheimer's Disease**. Lilly has dedicated over 30 years to supporting research efforts in Alzheimer's¹ and is committed to being part of a better future for people living with this devastating disease and their loved ones.

A GROWING GLOBAL **HEALTH CRISIS**

EVERY 3 SECONDS

someone around the world develops dementia primarily as a result of Alzheimer's disease²

55 MILLION PEOPLE with dementia expected to be ~140 million by 2050°

PEOPI F worldwide at risk of

developing dementia caused by Alzheimer's

Annually dementia costs healthcare systems

~S1 TRILLION by 2030 expected to reach >\$2.8 TRILLION³

AFFECTING INDIVIDUALS AND FAMILIES

On average caregivers spend **T** HOURS looking after a loved one with dementia,3 which can lead caregivers to experience:



-SUBSTANTIAL

-LOSS OF EARNINGS⁵

FINANCIAL COSTS⁵ -INCREASED ANXIETY



AND DEPRESSION5 - OTHER HEALTH ISSUES⁵ From the early stages of cognitive impairment, **ALZHEIMER'S LEADS TO SEVERE EMOTIONAL, SOCIAL, PSYCHOLOGICAL** & PRACTICAL **CHALLENGES** reducing life expectancy and independence^{6,7}



-ROUGHLY \$500 BILLIONis attributed to informal care³

Half the global cost of dementia

diagnosing and treating Alzheimer's disease earlier[®] **OBTAINING A TIMELY** TODAY

Preventing or delaying cognitive decline relies on



Access to ADVANCED

DIAGNOSTIC



occurs early enough for emerging therapies to be an option¹⁰

DIAGNOSIS CAN: Enable earlier access to treatment and support,

AND ACCURATE



Help those affected plan

for the future and maintain

independence for longer^{11,12}

their quality of life and

and better management of

co-morbid conditions¹¹



Offer people the chance to participate in clinical trials 11,12

Changing the course of Alzheimer's disease in the early stages can limit the damage caused and

reduce the impact on individuals, loved ones and

Encourage more people to recognise the early signs and seek help

https://doi.org/10.1002/gps.1652.

Alzheimer Dis Assoc Disord. 2009 Oct-Dec;23(4):306-14.

health services.

Lilly is calling for policies which:

Arm clinicians with the

tools to diagnose

earlier and more

accurately

Address gaps in care pathways

to improve

patient outcomes

GM-31058

References

- 1. Eli Lilly and Company. Alzheimer's webpage. Available at: https://www.lilly.com/disease-areas/alzheimers. Last accessed August 2022 Alzheimer's Disease International. Dementia Statistics. Available at: https://www.alzint.org/about/dementia-facts-figures/dementia-statistics/. Last accessed June 2022.
- Gustavsson, A, Norton, N, Fast, T, et al. Global estimates on the number of persons across the Alzheimer's disease continuum. Alzheimer's Dement. 2022; 1-13. https://doi.org/10.1002/alz.12694. 5. Lindeza P, et al. BMJ Supportive & Palliative Care 2020;0:1–12. doi:10.1136/bmjspcare-2020-002242.

3. World Health Organization. Dementia. Available at: https://www.who.int/news-room/fact-sheets/detail/dementia. Last accessed June 2022.

- 6. Mayo Clinic. Alzheimer's disease. Available at: https://www.mayoclinic.org/diseases-conditions/alzheimers-disease/in-depth/alzheimers-stages/art-20048448. 7. Abbas A, Monfared T, Byrnes MJ. The Humanistic and Economic Burden of Alzheimer's Disease. Neurol Ther. 2002. doi.org/10.1007/s40120-022-00335-x 8. Biagioni MC, Galvin JE. Using biomarkers to improve detection of Alzheimer's disease. Neurodegener Dis Manag. 2011 Apr;1(2):127-139.
- 11. Rasmussen, J., & Langerman, H. (2019). Alzheimer's Disease Why We Need Early Diagnosis. Degenerative neurological and neuromuscular disease, 9, 123–130. https://doi.org/10.2147/DNND.S228939.

10. Kotagal V, et al. Factors Associated with Cognitive Evaluations in the United States. Neurology 2015;84[1]:64-71] 12. Bradford A, Kunik ME, Schulz P, Williams SP, Singh H. Missed and delayed diagnosis of dementia in primary care: prevalence and contributing factors.

9. Waldemar G. et al. Access to diagnostic evaluation and treatment for dementia in Europe. International Journal of Geriatric Psychiatry. 2007; 22, 1.